

5

Integrals



5.1

Areas and Distances

Integral

مساحة

مسافة



The Area Problem

The Area Problem

We begin by attempting to solve the area problem: Find the area of the region S that lies under the curve $y = f(x)$ from a to b .

This means that S , illustrated in Figure 1, is bounded by the graph of a continuous function f [where $f(x) \geq 0$], the vertical lines $x = a$ and $x = b$, and the x -axis.




Figure 1

The Area Problem

مساحة

مساحة
For a rectangle, the area is defined as the product of the length and the width. The area of a triangle is half the base times the height.

مساحة

The area of a polygon is found by dividing it into triangles (as in Figure 2) and adding the areas of the triangles.




Figure 2

مساحة: طول \times عرض

مساحة قاعدة \times ارتفاع h

The Area Problem

However, it isn't so easy to find the area of a region with curved sides. We all have an intuitive idea of what the area of a region is. But part of the area problem is to make this intuitive idea precise by giving an exact definition of area.

Recall that in defining a tangent we first approximated the slope of the tangent line by slopes of secant lines and then we took the limit of these approximations.


We pursue a similar idea for areas. We first approximate the region S by rectangles and then we take the limit of the areas of these rectangles as we increase the number of rectangles.

Example 1

مثال

Use rectangles to estimate the area under the parabola $y = x^2$ from 0 to 1 (the parabolic region S illustrated in Figure 3).

استخدام
المثلثات
لتقدير المساحة




parabola

Figure 3

Example 1 – Solution

We first notice that the area of S must be somewhere between 0 and 1 because S is contained in a square with side length 1, but we can certainly do better than that.

Suppose we divide S into four strips $S_1, S_2, S_3,$ and S_4 by drawing the vertical lines $x = \frac{1}{4}, x = \frac{1}{2},$ and $x = \frac{3}{4}$ as in Figure 4(a).





$$0 + \frac{1}{n} \quad \frac{1}{4} + \frac{1}{n} \quad \boxed{\frac{2}{5}} + \frac{1}{n} \quad \frac{3}{5} + \frac{1}{n} \quad \downarrow$$

$$n \rightarrow \frac{\dot{1} - \dot{0}}{5} = \boxed{\frac{1}{5}}$$

$$\boxed{\frac{2}{5}} + \frac{1}{n} \downarrow \frac{1}{2}$$

$$\Delta x = \frac{\hat{a}(n) - \hat{a}(1)}{n}$$

موتة التقتيم

Example 1 – Solution

We can approximate each strip by a rectangle that has the same base as the strip and whose height is the same as the right edge of the strip [see Figure 4(b)].

In other words, the heights of these rectangles are the values of the function $f(x) = x^2$ at the *right endpoints* of the subintervals

$[0, \frac{1}{4}]$, $[\frac{1}{4}, \frac{1}{2}]$, $[\frac{1}{2}, \frac{3}{4}]$, and $[\frac{3}{4}, 1]$.

Each rectangle has width $\frac{1}{4}$ and the heights are $(\frac{1}{4})^2$, $(\frac{1}{2})^2$, $(\frac{3}{4})^2$, and 1^2 .




Figure 4(b)

Right end point
small

$$y = x^2$$

$$(1, 1)^2$$



$$(1, 1)^2$$

$\underbrace{1/4}_{1/4} \times d_{\text{area}}^n = \underline{\underline{\text{jumlah luas}}}$

[1]

$$\left(\frac{1}{u}\right)^2 = \text{الارتفاع}^2 \text{ طول}^2$$

[2]

$$\left(\frac{2}{u}\right)^2 = \text{الطول}^2$$

[3]

$$\left(\frac{3}{u}\right)^2 = \text{طول}^2$$

[4]

$$(1)^2 = \text{طول}^2$$

$$\text{Area} = \boxed{1} + \boxed{2} + \boxed{3} + \boxed{4}$$

$$\begin{aligned} \text{Area} &= \left(\frac{1}{u}\right) \left(\frac{1}{u}\right)^2 + \left(\frac{1}{u}\right) \left(\frac{2}{u}\right)^2 + \left(\frac{1}{u}\right) \left(\frac{3}{u}\right)^2 \\ &\quad + \left(\frac{1}{u}\right) (1)^2 \end{aligned}$$

$$\text{Area} = \frac{15}{32} = 0.46875$$

Example 1 – Solution


cont'd

If we let R_4 be the sum of the areas of these approximating rectangles, we get

$$R_4 = \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 + \frac{1}{4} \cdot 1^2$$

$$= \frac{15}{32}$$

$$= 0.46875$$



From Figure 4(b) we see that the area A of S is less than R_4 , so

$$A < 0.46875$$

Right end point

• دائماً ϵ \bar{U} نبدأ!

Example 1 – Solution

cont'd

Instead of using the rectangles in Figure 4(b) we could use the smaller rectangles in Figure 5 whose heights are the values of f at the left endpoints of the subintervals. (The leftmost rectangle has collapsed because its height is 0.)




Figure 4(b)

Right endpoint





Figure 5

left end point



$$(0)^2 = 0$$

$$\underline{\underline{\left(\frac{1}{4}\right)^2}}$$

$$\left(\frac{1}{2}\right)^2 = \underline{\underline{\left(\frac{2}{4}\right)^2}}$$

$$\underline{\underline{\left(\frac{3}{4}\right)^2}}$$

1] الارتفاع .

2] الارتفاع .

3] الارتفاع .

4] الارتفاع .


مساحة مستطيل \Rightarrow طول \times العرض .

Example 1 – Solution

cont'd

The sum of the areas of these approximating rectangles is

$$\begin{aligned} L_4 &= \frac{1}{4} \cdot \underline{0^2} + \frac{1}{4} \cdot \underline{\left(\frac{1}{4}\right)^2} + \frac{1}{4} \cdot \underline{\left(\frac{1}{2}\right)^2} + \frac{1}{4} \cdot \underline{\left(\frac{3}{4}\right)^2} \\ &= \frac{7}{32} \\ &= 0.21875 \end{aligned}$$



We see that the area of S is larger than L_4 , so we lower and upper estimates for A :

$$\underline{0.21875} < A < \underline{0.46875}$$

We can repeat this procedure with a larger number of strips.

Example 1 – Solution

cont'd

Instead of using the rectangles in Figure 4(b) we could use the smaller rectangles in Figure 5 whose heights are the values of f at the left endpoints of the subintervals. (The leftmost rectangle has collapsed because its height is 0.)




Figure 4(b)

Right





Figure 5

Left


Example 1 – Solution

cont'd

Figure 6 shows what happens when we divide the region S into eight strips of equal width.



(a) Using left endpoints



(b) Using right endpoints

تزيان حلت

Approximating S with eight rectangles

Figure 6

Example 1 – Solution

cont'd

By computing the sum of the areas of the smaller rectangles (L_8) and the sum of the areas of the larger rectangles (R_8), we obtain better lower and upper estimates for A :

$$\underline{\underline{\text{نِسْبَة}}}$$
$$0.2734375 < A < 0.3984375$$
$$\underline{\underline{\text{نِسْبَة}}}$$

So one possible answer to the question is to say that the true area of S lies somewhere between 0.2734375 and 0.3984375.

We could obtain better estimates by increasing the number of strips.

Example 1 – Solution

cont'd

The table at the right shows the results of similar calculations (with a computer) using n rectangles whose heights are found with left endpoints (L_n) or right endpoints (R_n).

n	L_n	R_n
10	0.2850000	0.3850000
20	0.3087500	0.3587500
30	0.3168519	0.3501852
50	0.3234000	0.3434000
100	0.3283500	0.3383500
1000	0.3328335	0.3338335

left *Right*

الف صغیر →


≈ 0,33

In particular, we see by using 50 strips that the area lies between 0.3234 and 0.3434. With 1000 strips we narrow it down even more: A lies between 0.3328335 and 0.3338335.

A good estimate is obtained by averaging these numbers:
 $A \approx 0.3333335$.

The Area Problem

From Figures 8 and 9 it appears that, as n increases, both L_n and R_n become better and better approximations to the area of S .




Right endpoints produce upper sums because $f(x) = x^2$ is increasing.

Figure 8

كل ما قسمت اكثر كود كل اقل .

The Area Problem



Left endpoints produce lower sums because $f(x) = x^2$ is increasing.

Figure 9

Therefore we *define* the area A to be the limit of the sums of the areas of the approximating rectangles, that is,

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} L_n = \boxed{\frac{1}{3}}$$

$$\frac{1}{3} \approx 0.3333 \dots$$

Handwritten notes:
 - The limit symbol $\lim_{n \rightarrow \infty}$ is circled in purple.
 - The fraction $\frac{1}{3}$ is boxed in purple.
 - The decimal expansion $0.3333 \dots$ is written in purple.
 - There are additional purple scribbles and arrows on the left side of the equation.

The Area Problem

We start by subdividing S into n strips S_1, S_2, \dots, S_n of equal width as in Figure 10.




Figure 10

The width of the interval $[a, b]$ is $b - a$, so the width of each of the n strips is

$$\Delta x = \frac{b - a}{n}$$

The Area Problem

التقسيم بالتساوي

Therefore we define the area A of the region S in the following way.

المساحة = مجموع المساحات = $f(\Delta x) \Delta x$

2 Definition The **area** A of the region S that lies under the graph of the continuous function f is the limit of the sum of the areas of approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1) \Delta x + f(x_2) \Delta x + \cdots + f(x_n) \Delta x]$$

It can be proved that the limit in Definition 2 always exists, since we are assuming that f is continuous. It can also be shown that we get the same value if we use left endpoints:

3
$$A = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} [f(x_0) \Delta x + f(x_1) \Delta x + \cdots + f(x_{n-1}) \Delta x]$$

Right | left $\rightarrow A(\Delta x)$

The Area Problem

In fact, instead of using left endpoints or right endpoints, we could take the height of the i th rectangle to be the value of f at *any* number x_i^* in the i th subinterval $[x_{i-1}, x_i]$. We call the numbers $x_1^*, x_2^*, \dots, x_n^*$ the **sample points**.

Figure 13 shows approximating rectangles when the sample points are not chosen to be endpoints.




Figure 13

So a more general expression for the area of S is

4

$$A = \lim_{n \rightarrow \infty} [f(x_1^*) \Delta x + f(x_2^*) \Delta x + \cdots + f(x_n^*) \Delta x]$$

The Area Problem

Note:


It can be shown that an equivalent definition of area is the following: *A is the unique number that is smaller than all the upper sums and bigger than all the lower sums.*

We saw in Example 1, for instance, that the area ($A = \frac{1}{3}$) is trapped between all the left approximating sums L_n and all the right approximating sums R_n .

The function in the example, $f(x) = x^2$, happens to be increasing on $[0, 1]$ and so the lower sums arise from left endpoints and the upper sums from right endpoints.

The Area Problem

In general, we form **lower** (and **upper**) **sums** by choosing the sample points x_i^* so that $f(x_i^*)$ is the minimum (and maximum) value of f on the i th subinterval. (See Figure 14)



Lower sums (short rectangles) and upper sums (tall rectangles)

Figure 14

The Area Problem

$\sum \Rightarrow$ *مجموع*
 $1+2+\dots$

We often use **sigma notation** to write sums with many terms more compactly. For instance,

$$\sum_{i=1}^n f(x_i) \Delta x = f(x_1) \Delta x + f(x_2) \Delta x + \dots + f(x_n) \Delta x$$

Handwritten notes: The summation symbol and the entire equation are boxed. Under each term $f(x_i) \Delta x$, there are handwritten notes: "مساحة" (area) and "طول" (width). A wavy line connects the Δx terms across the equation.

So the expressions for area in Equations 2, 3, and 4 can be written as follows:

$\infty \Rightarrow$ *محدوده*
دراسته

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}) \Delta x$$



The Distance Problem

The Distance Problem

المسافة

Now let's consider the *distance problem*: Find the distance traveled by an object during a certain time period if the velocity of the object is known at all times.

If the velocity remains constant, then the distance problem is easy to solve by means of the formula

$$\text{مسافة} \quad \underline{\text{distance}} = \underline{\text{سرعة}} \times \underline{\text{الزمن}} \quad \text{velocity} \times \text{time}$$

But if the velocity varies, it's not so easy to find the distance traveled.

متغيرة

$$\frac{km}{h} \Rightarrow \frac{\vec{a} \text{ المسافة}}{\text{الزمن}} \neq \frac{\text{السرعة}}{1}$$

$$\text{الزمن} \times \text{السرعة} = \vec{a} \text{ المسافة}, c =$$

Example 4

Suppose the odometer on our car is broken and we want to estimate the distance driven over a 30-second time interval. We take speedometer readings every five seconds and record them in the following table:

Time (s)	0	5	10	15	20	25	30
Velocity (mi/h)	17	21	24	29	32	31	28

In order to have the time and the velocity in consistent units, let's convert the velocity readings to feet per second ($1 \text{ mi/h} = 5280/3600 \text{ ft/s}$):

Time (s)	0	5	10	15	20	25	30
Velocity (ft/s)	25	31	35	43	47	45	41

تفكيرية

$$\frac{1 \text{ mi}}{\text{h}} \Rightarrow \frac{5280}{3600} \frac{\text{ft}}{\text{s}}$$

$$17 \frac{\text{mi}}{\text{h}} = 17 \times \frac{5280}{3600} \frac{\text{ft}}{\text{s}} = 24,933 \approx \underline{\underline{25}}$$

$$\begin{aligned} & \overset{[1]}{(25 \times 5)} + \overset{[2]}{(31 \times 5)} + \overset{[3]}{(35 \times 5)} + \overset{[4]}{(43 \times 5)} + \overset{[5]}{(47 \times 5)} \\ & + \overset{[6]}{(45 \times 5)} \Rightarrow \underline{\underline{1130}} \frac{\text{ft}}{\text{s}} \quad \underline{\underline{1120}} \end{aligned}$$

Example 4

During the first five seconds the velocity doesn't change very much, so we can estimate the distance traveled during that time by assuming that the velocity is constant.

If we take the velocity during that time interval to be the initial velocity (25 ft/s), then we obtain the approximate distance traveled during the first five seconds:

$$25 \text{ ft/s} \times 5 \text{ s} = \boxed{125 \text{ ft}}$$

Example 4

If we add similar estimates for the other time intervals, we obtain an estimate for the total distance traveled:

$$(25 \times 5) + (31 \times 5) + (35 \times 5) + (43 \times 5) + (47 \times 5) + (45 \times 5) = 1130 \text{ ft}$$

We could just as well have used the velocity at the end of each time period instead of the velocity at the beginning as our assumed constant velocity.

Then our estimate becomes

$$(31 \times 5) + (35 \times 5) + (43 \times 5) + (47 \times 5) + (45 \times 5) + (41 \times 5) = 1210 \text{ ft}$$

If we had wanted a more accurate estimate, we could have taken velocity readings every two seconds, or even every second.

mit ✓
Right